

Noctuidae

Acronictinae

Apsarasa radians (Westwood, 1848)



Taxonomy:

Noctua radians Westwood, 1848: 58.–
India (Assam).

Apsarasa figurata Moore, 1877: 604.–
India (Andamans).

Apsarasa nigrotarsata Grünberg,
1911: 163.– .


Apsarasa dajakana Grünberg, 1911:
163.– Indonesia (Borneo).

Hostplant

Flight period: v.

Altitude: 340-900 m.

Imago



Distribution map

Noctuidae

Acronictinae

Diphthera pallida Moore, 1867



Taxonomy:

Diphthera pallida Moore, 1867: 46.—
India (Bengal).


Hostplant

Flight period: viii.

Altitude: 2700-2910 m.

Imago

Diphthera pallida Moore, 1867



Distribution map

Noctuidae

Amphiperinae

Callyna jugaria Walker, 1858



Imago


Taxonomy :

Callyna jugaria Walker, 1858: 1809.–

Hostplant:

Flight period: viii.

Altitude: 875 m.



Distribution map

Noctuidae

Amphiperinae

Callyna monoleuca Walker, 1858



Imago


Taxonomy :

Callyna monoleuca Walker, 1858:
1667.- .

Hostplant:

Flight period: v, viii.

Altitude: 875-1975 m.



Distribution map

Noctuidae

Bagisarinae

Xanthodes transversa Guenée, 1852



Taxonomy

Xanthodes transversa Guenée, 1852:
211.- .

Xanthodes migrator Walker, 1858:
779.- .


Trileuca dentalis Smith, 1891: 123.- .

Hostplant

Flight period: v.

Altitude: 340-1750 m.

Imago



Distribution map

Noctuidae

Dyopsinae

Moma champa Moore, 1879



Imago


Taxonomy :

Moma champa Moore, 1879: 403.–
India (Dharmasala)..

Hostplant

Flight period: viii.-ix.

Altitude: 2700-2910 m.



Distribution map

Noctuidae

Hadeninae

Tiracola plagiata (Walker, 1857)



Taxonomy:

Agrotis plagiata Walker, 1857: 740.–
Ceylon.

Agrotis plagifera Walker, 1857: 741.–
Ceylon.


Agrotis spectabilis Walker, 1865: 704.
– Australia (Queensland).

Hostplant

Flight period: v, ix.

Altitude: 1555-2910 m.

Imago



Distribution map

Noctuidae

Xyleninae

Trachea auriplena (Walker, 1857)



Imago

Taxonomy:


Eurois ? auriplena Walker, 1857: 557.–
Ceylon.

Hostplant

Flight period: viii-ix.

Altitude: 2275-2910 m.

Remarks: specs uncertain id.



Distribution map

Noctuidae

Noctuinae

Actinotia intermediata (Bremer, 1861)



Taxonomy:


Chloantha intermediata Bremer, 1861:
489.- Corea (?).

Hostplant:

Flight period: ix.

Altitude: 2910 m.

Imago



Distribution map

Noctuidae

Noctuinae

Xestia c-nigrum (Linnaeus, 1758)



Taxonomy


Phalaena (Noctua) c-nigrum Linnaeus, 1758: 516.– Europe.

Hostplant: *Chamaenerion angustifolium*, *Stellaria media*.

Flight period: vi, viii-ix.

Altitude: 2890-2925 m.

Imago



Distribution map

Noctuidae

Pantheinae

Donda eurychlora Walker, 1858



Taxonomy:


Donda eurychlora Walker, 1858: .- .

Hostplant:

Flight period: viii.

Altitude: 875 m.

Imago



Distribution map

Noctuidae

Pantheinae

Trisuloides sericea Butler, 1881




Imago

Taxonomy: *Trisuloides sericea* Butler, 1881: 36.– India (Sikkim).

Hostplant:

Flight period: viii.

Altitude: 875 m.



Distribution map

Noctuidae

Plusiinae

Chrysodeixis acuta (Walker, 1857)



Taxonomy:

Plusia acuta Walker, [1858]: 922.–
Congo.

Neoplusia furihatai Okano, 1963: 90.–
Japan (Honshu).


Shensiplusia nigribursa Chou & Lu,
1974: 68.– China (Shensi).

Hostplant

Flight period: viii-ix.

Altitude: 2750-2910 m.

Imago



Distribution map

Noctuidae

Pluosiinae

Trichoplusia orichalcea (Fabricius, 1775)



Taxonomy:

Noctua orichalcea Fabricius, 1775:
607.– Europe.

Phalaena chrysitina Martyn, 1797:
plate 21.– .


Noctua aurifera Hübner, [1813]: 463.-
Europe.

Hostplant:

Flight period: v, viii.

Altitude: 875-1975 m.

Imago



Distribution map

Noctuidae

Xyleninae

Karana gemmifera (Walker, 1858)



Taxonomy:

Plusia gemmifera Walker, [1858]: 934 .-


Kerana[sic] similis Moore, 1888: 407.-
India (Himachal Pradesh).

Hostplant:

Flight period: v.

Altitude: 2040 m.

Imago



Distribution map

Noctuidae

Xyleninae

Olivenebula pulcherrima Moore, 1867



Taxonomy:

Epilecta pulcherrima Moore, 1867: 54.
– India (Darjeeling).

Eliochroea opulenta Butler, 1889: 73.–
India (Dharmasala).


Polyphaenis largeteui Oberthür,
1881: 19.– China (Kouy-Tcheou).

Hostplant:

Flight period: ix.

Altitude: 2910 m.

Imago



Distribution map

Noctuidae

Xyleninae

Phlogophora albovittata (Moore, 1867)



Taxonomy:


Euplexia albovittata Moore, 1867: 57.–
India (Darjeeling).

Hostplant:

Flight period: ix

Altitude: 2910 m.

Imago



Distribution map

Noctuidae

Xyleninae

Phlogophora subpurpurea Leech, 1900



Imago


Taxonomy:

Phlogophora subpurpurea Leech,
1900: 71.– China (Sichuan).

Hostplant

Flight period: ix.

Altitude: 2910 m.



Distribution map